

Applications of the derivative operator on some subclasses of analytic functions

Ali Omar

Introduction

We begin by letting $\mathcal{U} = \{z \in \mathbb{C}: |z| < 1\}$ be the open unit disk of the complex plane and \mathcal{A} denote the class of functions of the form

$$f(z) = z + \sum_{k=2}^{\infty} a_k z^k,$$

which are analytic in \mathcal{U} and satisfy the following usual normalization conditions $f(0) = f'(0) - 1 = 0$.

Also, let S be the class of functions $f \in \mathcal{A}$ which are univalent in \mathcal{U} . Given two analytic functions f and g , the subordination between them is written as $f < g$ or $f(z) < g(z)$, ($z \in \mathcal{U}$). In addition, we say $f(z)$ is subordinate to $g(z)$ if there is a Schwarz function w with $w(z) = 0$, $|w(z)| < 1$, ($z \in \mathcal{U}$) such that $f(z) = g(w(z))$ for all $z \in \mathcal{U}$. Furthermore, if $g(z)$ is univalent in \mathcal{U} , then $f < g$ if and only if $f(0) = g(0)$ and $f(\mathcal{U}) \subseteq g(\mathcal{U})$.

In [6], [7], for a function $f \in \mathcal{A}$ and $0 < q < 1$ Jackson defined the q -derivative operator D_q as follows:

$$D_q f(z) = \frac{f(qz) - f(z)}{(q-1)z}, \quad (z \neq 0)$$

and $D_q f(0) = f'(0)$ and $D_q^2 f(z) = D_q(D_q f(z))$. In case $f(z) = z^k$ for k is a positive integer, the q -derivative of $f(z)$ is given by

$$D_q z^k = \frac{z^k - (zq)^k}{z(1-q)} = [k]_q z^{k-1},$$

where $[k]_q$ defined by

$$[k]_q = \frac{1 - q^k}{1 - q}$$

As $q \rightarrow 1^-$ and $k \in \mathbb{N}$, $[k]_q \rightarrow k$

From (1.1) and (1.2) we get that

$$D_q f(z) = 1 + \sum_{k=2}^{\infty} [k]_q a_k z^{k-1}.$$

As a right inverse, Jackson [6] introduced the q -integral

$$\int_0^z f(t) d_q t = z(1-q) \sum_{k=0}^{\infty} q^k f(zq^k)$$

The authors in [1] defined the q - analogue of Ruscheweyh Operator \mathcal{R}_q^λ by

$$\mathcal{R}_q^\lambda f(z) = z + \sum_{k=2}^{\infty} \frac{[k + \lambda - 1]_q!}{[\lambda]_q! [k - 1]_q!} a_k z^k$$

where $[k]_q!$ defined by :

$$[k]_q! = \begin{cases} [k]_q [k - 1]_q \dots [1]_q, & k = 1, 2, \dots \\ 1; & k = 0 \end{cases}$$

All the details about q - calculus used in this paper can be found in [3] and [4].

Also, as $q \rightarrow 1^-$ we have

$$\begin{aligned} \lim_{q \rightarrow 1^-} \mathcal{R}_q^\lambda f(z) &= z + \lim_{q \rightarrow 1^-} \left[\sum_{k=2}^{\infty} \frac{[k + \lambda - 1]_q!}{[\lambda]_q! [k - 1]_q!} a_k z^k \right] \\ &= z + \sum_{k=2}^{\infty} \frac{(k + \lambda - 1)!}{(\lambda)! (k - 1)!} a_k z^k \\ &= \mathcal{R}^\lambda f(z) \end{aligned}$$

where $\mathcal{R}^\lambda f(z)$ is Russcheweyh differential operator which was defined in [13] and has been studied by several authors, for example [9] and [15].

The following class of analytic functions can be defined as a result of full utilization of subordination and q -derivative principle.

Definition 1.1 For $\phi \in \mathcal{P}, b \in \mathbb{C}^* = \mathbb{C} \setminus \{0\}, \beta \geq 0$ and $\lambda > -1$. The function f is said to be in the class $\mathcal{M}_{q,b,\lambda}^\beta(\phi)$ if it satisfies the condition :

$$1 + \frac{1}{b} \left[\frac{z D_q (\mathcal{R}_q^\lambda f(z)) + \beta z^2 D_q (D_q (\mathcal{R}_q^\lambda f(z)))}{\mathcal{R}_q^\lambda f(z)} - 1 \right] < \phi(z)$$

Remark 1.1 It can be seen that, by specializing the parameters, the class $\mathcal{M}_{q,b,\lambda}^\beta(\phi)$ is reduced to numerous known subclasses of analytic functions, for example:

- If $b = 1$ and $\beta = 0$ we obtain $\mathcal{M}_{q,1,\lambda}^\beta(\phi) \equiv \mathcal{S}_{q,\lambda}^*(\phi)$ (Aldweby and Darus [2]). - If $\lambda = 0$ and $\beta = 0$ we obtain $\mathcal{M}_{q,1,\lambda}^\beta(\phi) \equiv \mathcal{S}_{q,b}(\phi)$ (Seoudy and Aouf [14]).
- If $\lambda = 0, \lim_{q \rightarrow 1} \mathcal{M}_{q,b,0}^\beta(\phi) = \mathcal{M}_{\alpha,b}(\phi)$ (Suchithra et al. [16]).
- If $\lambda = 0$ and $\beta = 0, \lim_{q \rightarrow 1} \mathcal{M}_{q,b,0}^0(\phi) = \mathcal{S}_b(\phi)$ (Ravichandran et al. [11]).

- If $\lambda = 0$ and $\beta = 0$, $\lim_{q \rightarrow 1} \mathcal{M}_{q,b,0}^0 \left(\frac{1+z}{1-z} \right) = \mathcal{S}^*(b)$ (Nasr and Aouf [10]).
- If $\lambda = 0$ and $\beta = 0$, $\lim_{q \rightarrow 1} \mathcal{M}_{q,1-\eta,0}^0 \left(\frac{1+z}{1-z} \right) = \mathcal{S}^*(\eta)$ (Robertson [12])

In 1933 a study conducted by Fekete and Szegő [5] revealed that the maximum value of $|a_3 - \mu a_2^2|$ as a function of the real parameters μ , for functions belonging to the class \mathcal{S} . Following this, several attempts and researchers solved the Fekete-Szegő problem for various classes of \mathcal{S} class in addition to subclasses of functions in \mathcal{A} . Similar work is shown in [2], [14] and [16]. In this paper, the Fekete-Szegő inequality is obtained for functions in a more general class of $\mathcal{M}_{q,b,\lambda}^\beta(\phi)$ for functions defined above.

In order to prove and validate our results, following preliminary results are required.

4. Preliminary Results

Lemma 2.1 [8] If $p(z) = 1 + c_1 z + c_2 z^2 + \dots \in \mathcal{P}$ of positive real part is in \mathcal{U} and μ a complex number, then

$$|c_2 - \mu c_1^2| \leq 2 \max\{1, |2\mu - 1|\}.$$

The result is sharp given by

$$p(z) = \frac{1+z}{1-z} \text{ and } p(z) = \frac{1+z^2}{1-z^2}.$$

Lemma 2.2 [8] If $p(z) = 1 + c_1 z + c_2 z^2 + \dots$ is a function with positive real part, then

$$|c_2 - v c_1^2| = \begin{cases} -4v + 2, & \text{if } v \leq 0 \\ 2, & \text{if } 0 \leq v \leq 1 \\ 4v - 2, & \text{if } v \geq 1 \end{cases}$$

5. Main Results

Now is our theorems using similar methods studied by Aldweby and Darus in [2].

Theorem 3.1 Let $\phi(z) = 1 + B_1 z + B_2 z^2 + \dots \in \mathcal{P}$. If f given by (1.1) is in the class $\mathcal{M}_{q,b,\lambda}^\beta(\phi)$ and μ is a complex number, then

$$|a_3 - \mu a_2^2| \leq \frac{[2]_q |b B_1|}{[(1 + [2]_q \beta)[3]_q - 1][\lambda + 1]_q [\lambda + 2]_q} \cdot \max \left\{ 1, \left| \frac{B_2}{B_1} + \frac{[2]_q [(1 + \beta)[2]_q - 1][\lambda + 1]_q - \mu [(1 + [2]_q \beta)[3]_q - 1][\lambda + 2]_q}{[2]_q [(1 + \beta)[2]_q - 1]^2 [\lambda + 1]_q} b B_1 \right| \right\}.$$

The result is sharp.

Proof. If $f \in \mathcal{M}_{q,b,\lambda}^\beta(\phi)$, then there is a function $w(z)$ in \mathcal{U} with $w(0) = 0$ and $|w(z)| < 1$ in \mathcal{U} such that

$$1 + \frac{1}{b} \left[\frac{zD_q(\mathcal{R}_q^\lambda f(z)) + \beta z^2 D_q(D_q(\mathcal{R}_q^\lambda f(z)))}{\mathcal{R}_q^\lambda f(z)} - 1 \right] = \phi(w(z)).$$

Define the function $p(z)$ by

$$p(z) = \frac{1 + w(z)}{1 - w(z)} = 1 + d_1 z + d_2 z^2 + \dots.$$

Since $w(z)$ is a Schwarz function, immediately $\operatorname{Re}(p(z)) > 0$ and $p(0) = 1$. Therefore,

$$\begin{aligned} \phi(w(z)) &= \phi\left(\frac{p(z) - 1}{p(z) + 1}\right) \\ &= \phi\left(\frac{1}{2} \left[d_1 z + \left(d_2 - \frac{d_1^2}{2}\right) z^2 + \left(d_3 - d_1 d_2 + \frac{d_1^3}{4}\right) z^3 + \dots \right] \right) \\ &= 1 + \frac{1}{2} B_1 d_1 z + \left[\frac{1}{2} B_1 \left(d_2 - \frac{d_1^2}{2}\right) + \frac{1}{4} B_2 d_1^2 \right] z^2 + \dots \end{aligned}$$

since

$$\begin{aligned} 1 + \frac{1}{b} \left[\frac{zD_q(\mathcal{R}_q^\lambda f(z)) + \beta z^2 D_q(D_q(\mathcal{R}_q^\lambda f(z)))}{\mathcal{R}_q^\lambda f(z)} - 1 \right] &= 1 + \frac{[(1 + \beta)[2]_q - 1][\lambda + 1]_q}{b} a_2 z \\ &+ \left[\frac{[(1 + [2]_q \beta)[3]_q - 1][\lambda + 1]_q [\lambda + 2]_q}{[2]_q b} a_3 - \frac{[(1 + \beta)[2]_q - 1][\lambda + 1]_q^2}{b} a_2^2 \right] z^2 + \dots \end{aligned}$$

From (3.2) and (3.3), we obtain

$$\begin{aligned} \frac{[(1 + \beta)[2]_q - 1][\lambda + 1]_q}{b} a_2 &= \frac{B_1 d_1}{2} \\ \frac{[(1 + [2]_q \beta)[3]_q - 1][\lambda + 1]_q [\lambda + 2]_q}{[2]_q b} a_3 - \frac{[(1 + \beta)[2]_q - 1][\lambda + 1]_q^2}{b} a_2^2 \\ &= \frac{1}{2} B_1 \left(d_2 - \frac{d_1^2}{2}\right) + \frac{1}{4} B_2 d_1^2, \end{aligned}$$

or, equivalently,

$$\begin{aligned}
a_2 &= \frac{bB_1d_1}{2[(1+\beta)[2]_q-1][\lambda+1]_q}, \\
&= \frac{[2]_qbB_1d_2}{2[(1+[2]_q\beta)[3]_q-1][\lambda+1]_q[\lambda+2]_q} + \\
&\quad \frac{[2]_qd_1^2}{4[(1+[2]_q\beta)[3]_q-1][\lambda+1]_q[\lambda+2]_q} \left[\frac{b^2B_1^2}{[(1+\beta)[2]_q-1]} - b(B_1-B_2) \right].
\end{aligned}$$

Therefore,

$$a_3 - \mu a_2^2 = \frac{[2]_qbB_1}{2[(1+[2]_q\beta)[3]_q-1][\lambda+1]_q[\lambda+2]_q} (d_2 - \nu d_1^2)$$

where

$$\nu = \frac{1}{2} \left[1 - \frac{B_2}{B_1} - \frac{b[2]_q[(1+\beta)[2]_q-1][\lambda+1]_q - \mu b[(1+[2]_q\beta)[3]_q-1][\lambda+2]_q}{[2]_q[(1+\beta)[2]_q-1]^2[\lambda+1]_q} B_1 \right].$$

Our result now follows by an application of Lemma 2.1. Again by Lemma 2.1, the equality in (3.1) is gained for

$$p(z) = \frac{1+z}{1-z} \quad \text{or} \quad p(z) = \frac{1+z^2}{1-z^2}.$$

Thus the proof of Theorem (3.1) is complete.

Setting $q \rightarrow 1^-$ and $\lambda = 0$ in Theorem 3.1, we obtain the following result

Corollary 3.1 [16] Let $\phi(z) = 1 + B_1z + B_2z^2 + \dots \in \mathcal{P}$. If f given by (1.1) is in the class $\mathcal{M}_{\beta,b}(\phi)$ and μ is a complex number, then

$$|a_3 - \mu a_2^2| \leq \frac{|B_1b|}{2(1+3\beta)} \max \left\{ 1, \left| \frac{B_2}{B_1} + \left[\frac{(1+2\beta) - 2\mu(1+3\beta)}{(1+2\beta)^2} \right] bB_1 \right| \right\}.$$

The result is sharp.

Taking $\lambda = 0$ and $\beta = 0$ in Theorem 3.1, we obtain the following result.

Corollary 3.2 [14] Let $\phi(z) = 1 + B_1z + B_2z^2 + \dots \in \mathcal{P}$. If f given by (1.1) is in the class $\mathcal{S}_{q,b}(\phi)$ and μ is a complex number, then

$$|a_3 - \mu a_2^2| \leq \frac{|B_1b|}{[3]_q-1} \max \left\{ 1, \left| \frac{B_2}{B_1} + \left[\frac{([2]_q-1) - \mu([3]_q-1)}{([2]_q-1)^2} \right] bB_1 \right| \right\}.$$

The result is sharp.

By using Lemma 2.2, we have the following theorem.

Theorem 3.2 Let $\phi(z) = 1 + B_1z + B_2z^2 + \dots$ with $B_1 > 0$ and $B_2 \geq 0$. Let

$$\begin{aligned}\rho_1 &= \frac{[2]_q[(1+\beta)[2]_q - 1]^2[\lambda+1]_q[bB_1^2 + [(1+\beta)[2]_q - 1](B_2 - B_1)]}{bB_1^2[\lambda+2]_q[(1+\beta)[2]_q - 1][(1+[2]_q\beta)[3]_q - 1]}, \\ \rho_2 &= \frac{[2]_q[(1+\beta)[2]_q - 1]^2[\lambda+1]_q[bB_1^2 + [(1+\beta)[2]_q - 1](B_2 + B_1)]}{bB_1^2[\lambda+2]_q[(1+\beta)[2]_q - 1][(1+[2]_q\beta)[3]_q - 1]}.\end{aligned}$$

Let f given by (1.1) be in the class $\mathcal{M}_{q,b,\lambda}^\beta(\phi)$. Then

$$|a_3 - \mu a_2^2| \leq \begin{cases} \frac{[2]_q b B_2}{\gamma} + \frac{[2]_q b B_1^2}{\gamma} \left(\frac{b[2]_q[(1+\beta)[2]_q - 1][\lambda+1]_q - \mu b \gamma / [\lambda+1]_q}{[2]_q[(1+\beta)[2]_q - 1]^2[\lambda+1]_q} \right), & \text{if } \mu \leq \rho_1; \\ \frac{[2]_q b B_1}{\gamma}, & \text{if } \rho_1 \leq \mu \leq \rho_2 \\ -\frac{[2]_q b B_2}{\gamma} - \frac{[2]_q b B_1^2}{\gamma} \left(\frac{b[2]_q[(1+\beta)[2]_q - 1][\lambda+1]_q - \mu b \gamma / [\lambda+1]_q}{[2]_q[(1+\beta)[2]_q - 1]^2[\lambda+1]_q} \right), & \text{if } \mu \geq \rho_2, \end{cases}$$

where $\gamma = [(1+[2]_q\beta)[3]_q - 1][\lambda+1]_q[\lambda+2]_q$.

Proof. First, let $\mu \leq \rho_1$

$$\begin{aligned}|a_3 - \mu a_2^2| &\leq \frac{[2]_q b B_1}{2[(1+[2]_q\beta)[3]_q - 1][\lambda+1]_q[\lambda+2]_q} [-4\nu + 2] \\ &\leq \frac{[2]_q b B_2}{\gamma} + \frac{[2]_q b B_1^2}{\gamma} \left(\frac{b[2]_q[(1+\beta)[2]_q - 1][\lambda+1]_q - \mu b \gamma / [\lambda+1]_q}{[2]_q[(1+\beta)[2]_q - 1]^2[\lambda+1]_q} \right).\end{aligned}$$

Now, let $\rho_1 \leq \mu \leq \rho_2$, then using the above calculation, we get

$$|a_3 - \mu a_2^2| \leq \frac{[2]_q b B_1}{[(1+[2]_q\beta)[3]_q - 1][\lambda+1]_q[\lambda+2]_q}.$$

Finally, if $\mu \geq \rho_2$, then

$$\begin{aligned}|a_3 - \mu a_2^2| &\leq \frac{[2]_q b B_1}{2[(1+[2]_q\beta)[3]_q - 1][\lambda+1]_q[\lambda+2]_q} [4\nu - 2] \\ &\leq -\frac{[2]_q b B_2}{\gamma} - \frac{[2]_q b B_1^2}{\gamma} \left(\frac{b[2]_q[(1+\beta)[2]_q - 1][\lambda+1]_q - \mu b \gamma / [\lambda+1]_q}{[2]_q[(1+\beta)[2]_q - 1]^2[\lambda+1]_q} \right),\end{aligned}$$

where $\gamma = [(1+[2]_q\beta)[3]_q - 1][\lambda+1]_q[\lambda+2]_q$.

Taking $\beta = 0$ in Theorem 3.2 and knowing that $[n]_q - 1 = q[n-1]_q$, $[n+1]_q = [n]_q + q^n$ and $[n+2]_q = [n]_q + q^n[2]_q$, we obtain the following result

Corollary 3.3 [2] Let $\varphi(z) = 1 + B_1 z + B_2 z^2 + \dots$ with $B_1 > 0$ and $B_2 \geq 0$. Let

$$\sigma_1 = \frac{([\lambda]_q + q^\lambda)(B_1^2 + q(B_2 - B_1))}{([\lambda]_q + q^\lambda[2]_q)B_1^2}, \sigma_2 = \frac{([\lambda]_q + q^\lambda)(B_1^2 + q(B_2 + B_1))}{([\lambda]_q + q^\lambda[2]_q)B_1^2}.$$

Let f given by (1.1) be in the class $\mathcal{S}_{q,\lambda}^*(\varphi)$. Then

$$|a_3 - \mu a_2^2| \leq \begin{cases} \frac{B_2}{q[\lambda+1]_q[\lambda+2]_q} + \frac{B_1^2}{q[\lambda+1]_q[\lambda+2]_q} \left(\frac{([\lambda]_q + q^\lambda - ([\lambda]_q + q^\lambda[2]_q)\mu}{q([\lambda]_q + q^\lambda)} \right), & \text{if } \mu \leq \sigma_1; \\ \frac{B_1}{q[\lambda+1]_q[\lambda+2]_q}, & \text{if } \sigma_1 \leq \mu \leq \sigma_2 \\ -\frac{B_2}{q[\lambda+1]_q[\lambda+2]_q} - \frac{B_1^2}{q[\lambda+1]_q[\lambda+2]_q} \left(\frac{([\lambda]_q + q^\lambda - ([\lambda]_q + q^\lambda[2]_q)\mu}{q([\lambda]_q + q^\lambda)} \right), & \text{if } \mu \geq \sigma_2. \end{cases}$$

6. References

1. M. Abramowitz and I. A. Stegun (Eds.), Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables, Dover Publications Inc., New York, 1965.
2. A. A. Attiya, Some applications of Mittag-Leffler function in the unit disk, Filomat 30(7) (2016), 2075-2081. <https://doi.org/10.2298/FIL1607075A>
3. D. Bansal and J. K. Prajapat, Certain geometric properties of the Mittag-Leffler functions, Complex Var. Elliptic Equ. 61(3) (2016), 338-350. <https://doi.org/10.1080/17476933.2015.1079628>
4. T. Bulboacă and G. Murugusundaramoorthy, Univalent functions with positive coefficients involving Pascal distribution series, Commun. Korean Math. Soc. 35(3) (2020), 867 – 877. <https://doi.org/10.4134/CKMS.c190413>
5. N. E. Cho, S. Y. Woo and S. Owa, Uniform convexity properties for hypergeometric functions, Fract. Calc. Appl. Anal. 5(3) (2002), 303-313.
6. P. L. Duren, Univalent Functions, Grundlehren der Mathematischen Wissenschaften Series 259, Springer Verlag, New York, 1983. [7] M. El-Deeb, T. Bulboacă and J. Dziok, Pascal distribution series connected with certain subclasses of univalent functions, Kyungpook Math. J. 59 (2019), 301-314. <https://doi.org/10.5666/KMJ.2019.59.2.301>

7. B. A. Frasin, An application of an operator associated with generalized Mittag-Leffler function, Konuralp J. Math. 7(1) (2019), 199-202.
8. B. A. Frasin, T. Al-Hawary and F. Yousef, Some properties of a linear operator involving generalized Mittag-Leffler function, Stud. Univ. Babeş-Bolyai Math. 65(1) (2020), 67 – 75. <https://doi.org/10.24193/subbmath.2020.1.06>
9. B. A. Frasin, T. Al-Hawary and F. Yousef, Necessary and sufficient conditions for hypergeometric functions to be in a subclass of analytic functions, Afr. Mat. 30(1-2) (2019), 223-230. <https://doi.org/10.1007/s13370-018-0638-5>
10. H. J. Haubold, A. M. Mathai and R. K. Saxena, Mittag-Leffler functions and their applications, J. Appl. Math. 2011(2011), Article ID 298628.
<https://doi.org/10.1155/2011/298628>
11. V. Kiryakova, Generalized Fractional Calculus and Applications, Pitman Research Notes in Mathematics Series 301, Longman Scientific & Technical, Harlow, John Wiley & Sons, Inc., New York, 1994.
12. E. Merkes and B. T. Scott, Starlike hypergeometric functions, Proc. Amer. Math. Soc. 12 (1961), 885-888.
13. G. M. Mittag-Leffler, Sur la nouvelle fonction $E(x)$, C. R. Acad. Sci. Paris 137(1903), 554 – 558.
14. G. Murugusundaramoorthy, Subordination results for spiral-like functions associated with the Srivastava-Attiya operator, Integral Transforms Spec. Funct. 23(2) (2012), 97-103. <https://doi.org/10.1080/10652469.2011.562501>
15. G. Murugusundaramoorthy, D. Răducanu and K. Vijaya, A class of spirallike functions defined by Ruscheweyh-type q -difference operator, Novi Sad J. Math. 49(2)(2019), 59 – 71. <https://doi.org/10.30755/NSJOM.08284>
16. G. Murugusundaramoorthy, K. Vijaya and S. Porwal, Some inclusion results of certain subclass of analytic functions associated with Poisson distribution series, Hacet. J. Math. Stat. 45(4) (2016), 1101-1107. <https://doi.org/10.15672/HJMS.20164513110>
17. G. Murugusundaramoorthy, Subclasses of starlike and convex functions involving Poisson distribution series, Afr. Mat. 28(2017), 1357-1366.
<https://doi.org/10.1007/s13370-017-0520-x>

18. S. Porwal and M. Kumar, A unified study on starlike and convex functions associated with Poisson distribution series, Afr. Mat. 27(5) (2016), 1021-1027.
<https://doi.org/10.1007/s13370-016-0398-z>
19. M. S. Robertson, On the theory of univalent functions, Ann. of Math. (2) 37(2) (1936), 374-408.
20. H. Silverman, Starlike and convexity properties for hypergeometric functions, J. Math. Anal. Appl. 172 (1993), 574-581. <https://doi.org/10.1006/jmaa.1993.1044>
21. H. M. Srivastava, G. Murugusundaramoorthy and S. Sivasubramanian, Hypergeometric functions in the parabolic starlike and uniformly convex domains, Integral Transforms Spec. Funct. 18 (2007), 511 – 520. <https://doi.org/10.1080/10652460701391324>
22. L. Späček, Contribution à la théorie des fonctions univalentes, Časopis Pro Pěstování Matematiky 62(1932), 12 – 19.
23. A. Swaminathan, Certain sufficient conditions on Gaussian hypergeometric functions, Journal of Inequalities in Pure and Applied Mathematics 5(4) (2004), Article ID 83, 10 pages.
24. A. Wiman, Über die nullstellen der funktionen $E_\alpha(x)$, Acta Math. 29 (1905), 217-134.
25. T. R. Prabhakar, A single integral equation with a generalized Mittag – Leffler function in the kernel, Yokohama Math. J. 19 (1971), 7 – 15.
26. Salah, J. and Darus, M., A note on Generalized Mittag-Leffler function and Application, Far East Journal of Mathematical Sciences (FJMS) 48 (1), 33–46, 2011.
27. [Omar Jawabreh](#), [Ahmad Abdel Qader](#), [Jamal Salah](#), [Khaled Al Mashrafi](#), [Emad Al Dein](#) [AL Fahmawee](#), [Basel J. A. Ali](#), Fractional Calculus Analysis of Tourism Mathematical Model, Progr. Fract. Differ. Appl. Vol. 9, No. S1, pp. 1 -11 202) DOI: 10.18576/pfda/09S101 URL: <https://www.naturalspublishing.com/Article.asp?ArtcID=26373>
28. Jamal Salah, Maryam Al Hashmi and Khaled Al Mashrafi, Some propositions on mathematical models of population growth, International Journal of Latest Trends in Engineering and Technology (IJLTET) Special Issue - ICTIMESH - 2022, 64-69.
URL: https://www.ijltet.org/journal_details.php?id=978&j_id=5050
29. Jamal Salah, Hameed Ur Rehman, and, Iman Al- Buwaiqi, Inclusion Results of a Generalized Mittag-Leffler-Type Poisson Distribution in the k-Uniformly Janowski

- Starlike and the k -Janowski Convex Functions, *Mathematics and Statistics*, Vol. 11, No. 1, pp. 22-27, 2023.
DOI: 10.13189/ms.2023.110103
30. Jamal Salah, Riemann and Big Bang Hypotheses, *NEUROQUANTOLOGY*, Vol. 20, No. 16, pp. 1592-1597, 2022. DOI: 10.48047/NQ.2022.20.16. NQ880156
 31. Jamal Salah, Hameed Ur Rehman, and, Iman Al- Buwaiqi, Subclasses of Spiral-Like Functions Associated with the Generalized Mittag-Leffler Function, *NEUROQUANTOLOGY*, Vol. 20, No. 15, pp. 4784-4799, 2022. DOI: 10.14704/NQ.2022.20.15. NQ88484
 32. Jamal Salah, Maryam Al Hashmi, Hameed Ur Rehman, Khaled Al Mashrafi, Modified Mathematical Models in Biology by the Means of Caputo Derivative of A Function with Respect to Another Exponential Function, *Mathematics and Statistics*, Vol. 10, No. 6, pp. 1194 - 1205, 2022. DOI: 10.13189/ms.2022.100605
 33. Jamal Salah, Hameed Ur Rehman, and, Iman Al- Buwaiqi, The Non-Trivial Zeros of the Riemann Zeta Function through Taylor Series Expansion and Incomplete Gamma Function, *Mathematics and Statistics*, vol. 10, no. 2, pp. 410-418, 2022, DOI: 10.13189/ms.2022.100216.
 34. Jamal Salah, Some Remarks and Propositions on Riemann Hypothesis, *Mathematics and Statistics*, vol. 9, no. 2, pp. 159-165, 2021, DOI: 10.13189/ms.2021.090210.
 35. Hameed Ur Rehman, Malina Darus, and Jamal Salah, Generalizing Certain Analytic Functions Correlative to the n -th Coefficient of Certain Class of Bi-Univalent Functions, *Hindawi Journal of Mathematics* Volume 2021, Article ID 6621315, 14 pages DOI: [10.1155/2021/6621315](https://doi.org/10.1155/2021/6621315)
 36. Salah, J., (2021). Applications of differential Subordination and Superordination Involving Certain Fractional Operator. *Current Topics on Mathematics and Computer Science*, vol. 1, pp. 7–16, 2021 DOI: [10.9734/bpi/ctmcs/v1/9050D](https://doi.org/10.9734/bpi/ctmcs/v1/9050D)
 37. Jamal Y. Mohammad Salah, The Consequence of the Analytic Continuity of Zeta Function Subject to an Additional Term and a Justification of the Location of the Non-Trivial Zeros, *International Journal of Science and Research (IJSR)*, Vol. 9, Issue 3, March 2020, pp. 1566-1569, March 2020, URL: https://www.ijsr.net/get_abstract.php?paper_id=SR20328201050
 38. Jamal Y. Mohammad Salah, An Alternative perspective to Riemann Hypothesis, *PSYCHOLOGY AND EDUCATION*, vol. 57, no. 9, pp. 1278-1281, 2020, URL: [file:///C:/Users/drjamals/Downloads/454-Article%20Text-738-1-10-20210128%20\(1\).pdf](file:///C:/Users/drjamals/Downloads/454-Article%20Text-738-1-10-20210128%20(1).pdf)
 39. Jamal Y. Mohammad Salah, Two Conditional proofs of Riemann Hypothesis, *International Journal of Sciences: Basic and Applied Research (IJSBAR)*, vol. 49, no. 1, pp. 74-83, 2020, URL: <https://gssrr.org/index.php/JournalOfBasicAndApplied/article/view/10720>
 40. Jamal Salah, TWO NEW EQUIVALENT STATEMENTS TO RIEMANN HYPOTHESIS, *Far East Journal of Mathematical Sciences (FJMS)* Volume 118, Issue1, 2019, Pages 1- 8. Sept 2019. DOI: [10.17654/MS118010001](https://doi.org/10.17654/MS118010001)
 41. Jamal Y. Salah, A NEW SUBCLASS OF UNIVALENT FUNCTIONS DEFINED BY THE MEANS OF JAMAL OPERATOR, *Far East Journal of Mathematical Sciences (FJMS)* Vol. 108, No. 2, 2018, pp. 389-399, 2018, Dec 2018, DOI: [10.17654/MS108020389](https://doi.org/10.17654/MS108020389).
 42. Hameed Ur Rehman, Maslina Darus and **Jamal Salah**, Graphing Examples of Starlike and Convex Functions of order β , *Appl. Math. Inf. Sci.* vol. 12, No. 3, pp. 509-515, 2018,

DOI:10.18576/amis/120305

43. Jamal Salah, Hameed Ur Rehman and Maslina Darus, A Note on Caputo's derivative interpretation in Economy, J. Appl. Math, pp. 1-7, 2018, DOI: 10.1155/2018/1260240.
44. Hameed Ur Rehman, Maslina Darus and Jamal Salah, Normalization of the generalized K– Mittag-Leffler function and ratio to its sequence of partial sums, BSG Proceedings, vol. 25, 2018, pp. 78-94.
45. Hameed Ur Rehman, Maslina Darus and Jamal Salah, COEFFICIENT PROPERTIES INVOLVING THE GENERALIZED K–MITTAG-LEFFLER FUNCTIONS, Trans. J. Math. Mecha. (TJMM) 9 (2017), No. 2, 155-164
46. Jamal Y. Salah. Closed-to-Convex Criterion Associated to the Modified Caputo's fractional Calculus Derivative Operator. Far East Journal of Mathematical Sciences (FJMS). Vol. 101, No. 1, pp. 55-59, 2017, DOI: 10.17654/MS101010055.
47. Jamal Y. Salah. Properties of the Modified Caputo's Derivative Operator for certain analytic functions. International Journal of Pure and Applied Mathematics. September 2016, Vol. 109, No. 3, pp. 665 – 671, 2014, DOI: 10.12732/ijpam. v109i3.14
48. Jamal Salah. A Note on the Modified Caputo's Derivative Operator. Far East Journal of Mathematical Sciences (FJMS). Vol. 100, No. 4, pp. 609-615, 2016,

DOI: 10.17654/MS100040609

49. S. Venkatesh, Jamal Salah, G. Sethuraman. "Some Results on E – Cordial Graphs", *International Journal of Mathematical Trends and Technology (IJMTT)*. V7:121-125 March 2014.
DOI: 10.14445/22315373/IJMTT-V7P516
50. Jamal Y. Salah, A Note on Gamma Function, International Journal of Modern Sciences and Engineering Technology (IJMSET), vol. 2, no. 8, pp. 58-64, 2015,
51. Jamal Y. Salah, On Riemann Hypothesis and Robin's Inequality, International Journal of Scientific and Innovative Mathematical Research (IJSIMR), vol. 3, no. 4, pp. 9-14, April 2015,
URL: <https://www.arcjournals.org/pdfs/ijsimr/v3-i4/4.pdf>
52. Jamal Y. Salah, **A Note on Riemann Zeta Function**, IOSR Journal of Engineering (IOSRJEN), vol. 06, no. 02, pp. 07-16, February 2016,
URL: [http://iosrjen.org/Papers/vol6_issue2%20\(part-3\)/B06230716.pdf](http://iosrjen.org/Papers/vol6_issue2%20(part-3)/B06230716.pdf)
53. Jamal Y. Salah, A NOTE ON THE HURWITZ ZETA FUNCTION, Far East Journal of Mathematical Sciences (FJMS), vol. 101, no. 12, pp. 2677-2683, June 2017,
DOI: [10.17654/MS101122677](https://doi.org/10.17654/MS101122677)
54. Jamal Salah and Venkatesh Srivastava. Inequalities on the Theory of Univalent Function. Journal of Mathematics and System Science, vol. 4, pp. 509-513, 2014.
DOI: 10.17265/2159-5291/2014.07.008
55. Jamal Salah. Neighborhood of a certain family of multivalent functions with negative coefficients. International Journal of Pure and Applied Mathematics. (IJPAM). Vol. 92, No 4, April 2014,
DOI: 10.12732/ijpam. v92i4.14
56. Jamal Salah. Fekete-Szego problems involving certain integral operator. International Journal of Mathematics Trends and Technology. IJMTT, vol. 7, no. 1, pp. 54-60, 2014.
DOI: 10.14445/22315373/IJMTT-V7P508
57. Jamal Salah. Subordination and superordination involving certain fractional operator. Asian Journal of Fuzzy and Applied Mathematics, vol. 1, pp. 98-107, 2013.

URL: <https://www.ajouronline.com/index.php/AJFAM/article/view/724>

58. Jamal Salah. Certain subclass of analytic functions associated with fractional calculus operator. *Trans. J. Math. Mecha*, vol. 3, no. 1, pp. 35-42, 2011.
URL: <http://tjmm.edyopress.ro/journal/11030106.pdf>
59. Jamal Salah. A note on Starlike functions of order α associated with a fractional calculus operator involving Caputo's fractional. *J. Appl. comp Sc. Math*, vol. 5, no. 1, pp. 97- 101, 2011,
URL: https://www.jacsm.ro/view/?pid=10_16
60. Jamal Salah and Maslina Darus. On convexity of general integral operators.
J. Anale. Universitatii. De Vest. Timisoara. Seria Matematica – Informatica XLIX. 1 (2011) 117-124.
61. Jamal Salah and Maslina Darus. A note on generalized Mittag-Leffler and Application. *Far East. Math Sc. (FJMS)*, vol. 48, no. 1, pp. 33-46, 2011.
URL: <http://www.pphmj.com/abstract/5478.htm>
62. Jamal Salah and Maslina Darus. A subclass of uniformly convex functions associated with fractional calculus operator involving Caputo's fractional differentiation. *Acta. Univ. APL*. Vol. 24, pp. 295-306, 2010.
63. D. G. Cantor, "Power series with integral coefficients," *Bulletin of the American Mathematical Society*, vol. 69 , no. 3, pp. 362 – 367, 1963.
64. R. Vein and P. Dale, "Determinants and Their Applications in Mathematical Physics," in *Applied Mathematical Sciences*, vol. 134, Springer, New York, 1999.
65. R. Wilson, "Determinantal criteria for meromorphic functions," *Proceedings of the London Mathematical Society*, vol. s3-4, no. 1, pp. 357-374, 1954.
66. C. Pommerenke, "On the coefficients and Hankel determinants of univalent functions," *Journal of the London Mathematical Society*, vol. s1-41, no. 1, pp. 111-122, 1966. [5] C. Pommerenke, "On the Hankel determinants of univalent functions," *Mathematika*, vol. 14, no. 1, pp. 108-112, 1967.
67. D. Bansal, "Upper bound of second Hankel determinant for a new class of analytic functions," *Applied Mathematics Letters*, vol. 26, no. 1, pp. 103-107, 2013.
68. M. M. Elhosh, "On the second Hankel determinant of univalent functions," *Bull. Malays. Math. Soc.*, vol. 9, no. 1, pp. 23-25, 1986.
69. J. W. Noonan and D. K. Thomas, "On the Hankel determinants of areally mean p-valent functions," *Proceedings of the London Mathematical Society*, vol. s3-25, no. 3, pp. 503-524, 1972.
70. J. W. Noonan, "Coefficient differences and Hankel determinants of areally mean p-valent functions," *Proceedings of American Mathematical Society*, vol. 46, pp. 29-37, 1974.
71. J. W. Noonan and D. K. Thomas, "On the second Hankel determinant of areally mean p-valent functions," *Transactions of the American Mathematical Society*, vol. 223, pp. 337-346, 1976.
72. M. M. Elhosh, "On the second Hankel determinant of close-toconvex functions," *Bulletin of the Malaysian Mathematical Sciences Society*, vol. 9, no. 2, pp. 67-68, 1986.

73. M. Arif, K. I. Noor, and M. Raza, "Hankel determinant problem of a subclass of analytic functions," *Journal of Inequalities and Applications*, vol. 2012, Article ID 22, 2012
74. T. Hayami and S. Owa, "Generalized Hankel determinant for certain classes," *International Journal of Mathematics and Analysis*, vol. 4, no. 49-52, pp. 2573-2585, 2010.
75. T. Hayami and S. Owa, "Applications of Hankel determinant for p-valently starlike and convex functions of order α ," *Far East Journal of Applied Mathematics*, vol. 46, no. 1, pp. 1-23, 2010.
76. T. Hayami and S. Owa, "Hankel determinant for p-valently starlike and convex functions of order α ," *General Math*, vol. 17, no. 4, pp. 29-44, 2009.
77. A. Janteng, S. A. Halim, and M. Darus, "Hankel determinant for starlike and convex functions," *International Journal of Mathematics and Analysis*, vol. 1, no. 13-16, pp. 619-625, 2007.
78. A. K. Mishra and P. Gochhayat, "Second Hankel determinant for a class of analytic functions defined by fractional derivative," *International Journal of Mathematics and Mathematical Sciences*, vol. 2008, Article ID 153280, 10 pages, 2008.
79. N. Mohamed, D. Mohamad, and S. Cik Soh, "Second Hankel determinant for certain generalized classes of analytic functions," *International Journal of Mathematics and Analysis*, vol. 6, no. 17-20, pp. 807-812, 2012.
80. G. Murugusundaramoorthy and N. Magesh, "Coefficient inequalities for certain classes of analytic functions associated with Hankel determinant," *Bulletin of Mathematical Analysis and Applications*, vol. 1, no. 3, pp. 85-89, 2009.
81. K. I. Noor, "Higher order close-to-convex functions," *Japanese Journal of Mathematics*, vol. 37, no. 1, pp. 1-8, 1992.
82. K. I. Noor, "On the Hankel determinant problem for strongly close-to-convex functions," *Journal of Natural Geometry*, vol. 11, no. 1, pp. 29-34, 1997.
83. K. I. Noor, "On certain analytic functions related with strongly close-to-convex functions," *Applied Mathematics and Computation*, vol. 197, no. 1, pp. 149-157, 2008. [23] K. I. Noor and S. A. Al-Bany, "On Bazilevic functions," *International Journal of Mathematics and Mathematical Sciences*, vol. 10, no. 1, 88 pages, 1987.
84. K. I. Noor, "On analytic functions related with functions of bounded boundary rotation," *Commentarii Mathematici Universitatis Sancti Pauli*, vol. 30, no. 2, pp. 113-118, 1981.
85. K. I. Noor, "On meromorphic functions of bounded boundary rotation," *Caribb. J. Math.*, vol. 1, no. 3, pp. 95-103, 1982.
86. K. I. Noor, "Hankel determinant problem for the class of functions with bounded boundary rotation," *Revue Roumaine de Mathématiques Pures et Appliquées*, vol. 28, no. 8, pp. 731-739, 1983.
87. K. I. Noor and I. M. A. Al-Naggar, "On the Hankel determinant problem," *Journal of Natural Geometry*, vol. 14, no. 2, pp. 133-140, 1998.
88. R. M. Ali, S. K. Lee, V. Ravichandran, and S. Supramaniam, "The Fekete-Szegő coefficient functional for transforms of analytic functions," *Bulletin of the Iranian Mathematical Society*, vol. 35, no. 2, pp. 119-142, 2009.
89. R. M. Ali, V. Ravichandran, and N. Seenivasagan, "Coefficient bounds for p₋-valent functions," *Applied Mathematics and Computation*, vol. 187, no. 1, pp. 35-46, 2007. W. C. Ma and D. Minda, "A united treatment of some special classes of univalent functions," in *Proceedings of the Conference on Complex Analysis*, Tianjin, 1992.
90. R. K. Raina and J. Sokół, "On coefficient estimates for a certain class of starlike functions, Hacet," *Journal of Mathematics and Statistics*, vol. 44, no. 6, pp. 1427-1433, 2015.

91. G. Murugusundaramoorthy and T. Bulboacă, "Hankel determinants for new subclasses of analytic functions related to a shell shaped region," *Mathematics*, vol. 8, no. 6, article 1041, 2020.
92. S. Gandhi, "Radius Estimates for Three Leaf Function and Convex Combination of Starlike Functions," in *Mathematical Analysis I: Approximation Theory. ICRAPAM 2018*, vol. 306 of *Springer Proceedings in Mathematics & Statistics*, Springer, Singapore.
93. S. N. Malik, S. Mahmood, M. Raza, S. Farman, and S. Zainab, "Coefficient inequalities of functions associated with petal type domains," *Mathematics*, vol. 6, no. 12, p. 298, 2018.
94. K. Sharma, N. K. Jain, and V. Ravichandran, "Starlike functions associated with a cardioid," *Afrika Matematika*, vol. 27, pp. 923-939, 2016.
95. V. S. Masih and S. Kanas, "Subclasses of starlike and convex functions associated with the Limaçon domain," *Symmetry*, vol. 12, no. 6, p. 942, 2020.
96. Y. Yuzaimi, A. H. Suzeini, and A. B. Akbarally, "Subclass of starlike functions associated with a limaçon," *AIP Conference Proceedings*, vol. 1974, article 030023, 2018.
97. P. L. Duren, *Univalent Functions. Grundlehren der Mathematischen Wissenschaften*, vol. 259, Springer, New York, 1983.
98. U. Grenander and G. Szegő, *Toeplitz Forms and Their Applications*. California Monographs in Mathematical Sciences, University of California Press, Berkeley, 1958.
99. M. Abramowitz and I. A. Stegun (Eds.), *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, Dover Publications Inc., New York, 1965.
100. A. A. Attiya, Some applications of Mittag-Leffler function in the unit disk, *Filomat* 30(7) (2016), 2075-2081. <https://doi.org/10.2298/FIL1607075A>
101. D. Bansal and J. K. Prajapat, Certain geometric properties of the Mittag-Leffler functions, *Complex Var. Elliptic Equ.* 61(3) (2016), 338-350. <https://doi.org/10.1080/17476933.2015.1079628>
102. T. Bulboacă and G. Murugusundaramoorthy, Univalent functions with positive coefficients involving Pascal distribution series, *Commun. Korean Math. Soc.* 35(3) (2020), 867 – 877. <https://doi.org/10.4134/CKMS.c190413>
103. N. E. Cho, S. Y. Woo and S. Owa, Uniform convexity properties for hypergeometric functions, *Fract. Calc. Appl. Anal.* 5(3) (2002), 303-313.
104. P. L. Duren, *Univalent Functions, Grundlehren der Mathematischen Wissenschaften Series 259*, Springer Verlag, New York, 1983. [7] M. El-Deeb, T. Bulboacă and J. Dziok, Pascal distribution series connected with certain subclasses of univalent functions, *Kyungpook Math. J.* **59** (2019), 301-314. <https://doi.org/10.5666/KMJ.2019.59.2.301>
105. B. A. Frasin, An application of an operator associated with generalized Mittag-Leffler function, *Konuralp J. Math.* 7(1) (2019), 199-202.
106. B. A. Frasin, T. Al-Hawary and F. Yousef, Some properties of a linear operator involving generalized Mittag-Leffler function, *Stud. Univ. Babeş-Bolyai Math.* 65(1) (2020), 67 – 75. <https://doi.org/10.24193/subbbmath.2020.1.06>
107. B. A. Frasin, T. Al-Hawary and F. Yousef, Necessary and sufficient conditions for hypergeometric functions to be in a subclass of analytic functions, *Afr. Mat.* 30(1-2) (2019), 223-230. <https://doi.org/10.1007/s13370-018-0638-5>
108. H. J. Haubold, A. M. Mathai and R. K. Saxena, Mittag-Leffler functions and their applications, *J. Appl. Math.* 2011(2011), Article ID 298628. <https://doi.org/10.1155/2011/298628>
109. V. Kiryakova, *Generalized Fractional Calculus and Applications*, Pitman Research Notes in Mathematics Series 301, Longman Scientific & Technical, Harlow, John Wiley & Sons, Inc., New York, 1994.

110. E. Merkes and B. T. Scott, Starlike hypergeometric functions, *Proc. Amer. Math. Soc.* 12 (1961), 885-888.
111. G. M. Mittag-Leffler, Sur la nouvelle fonction $E(x)$, *C. R. Acad. Sci. Paris* 137(1903), 554 – 558.
112. G. Murugusundaramoorthy, Subordination results for spiral-like functions associated with the Srivastava-Attiya operator, *Integral Transforms Spec. Funct.* 23(2) (2012), 97-103. <https://doi.org/10.1080/10652469.2011.562501>
113. G. Murugusundaramoorthy, D. Răducanu and K. Vijaya, A class of spirallike functions defined by Ruscheweyh-type q -difference operator, *Novi Sad J. Math.* 49(2)(2019), 59 – 71. <https://doi.org/10.30755/NSJOM.08284>
114. G. Murugusundaramoorthy, K. Vijaya and S. Porwal, Some inclusion results of certain subclass of analytic functions associated with Poisson distribution series, *Hacet. J. Math. Stat.* 45(4) (2016), 1101-1107. <https://doi.org/10.15672/HJMS.20164513110>
115. G. Murugusundaramoorthy, Subclasses of starlike and convex functions involving Poisson distribution series, *Afr. Mat.* 28(2017), 1357-1366. <https://doi.org/10.1007/s13370-017-0520-x>
116. S. Porwal and M. Kumar, A unified study on starlike and convex functions associated with Poisson distribution series, *Afr. Mat.* 27(5) (2016), 1021-1027. <https://doi.org/10.1007/s13370-016-0398-z>
117. M. S. Robertson, On the theory of univalent functions, *Ann. of Math.* (2) 37(2) (1936), 374-408.
118. Salah, Jamal, Hameed Ur Rehman, and Iman Al-Buwaiqi. "The Non-Trivial Zeros of the Riemann Zeta Function through Taylor Series Expansion and Incomplete Gamma Function." *Mathematics and Statistics* 10.2 (2022): 410-418.
119. Rehman, Hameed Ur, Maslina Darus, and Jamal Salah. "Graphing Examples of Starlike and Convex Functions of order β ." *Appl. Math. Inf. Sci* 12.3 (2018): 509-515.
120. Salah, Jamal Y., and O. M. A. N. Ibra. "Properties of the Modified Caputo's Derivative Operator for certain analytic functions." *International Journal of Pure and Applied Mathematics* 109.3 (2014): 665-671.
121. Jamal Salah and Maslina Darus, On convexity of the general integral operators, *An. Univ. Vest Timis. Ser. Mat. -Inform.* 49(1) (2011), 117-124.
122. Salah, Jamal. "TWO NEW EQUIVALENT STATEMENTS TO RIEMANN HYPOTHESIS." (2019).
123. H. Silverman, Starlike and convexity properties for hypergeometric functions, *J. Math. Anal. Appl.* 172 (1993), 574-581. <https://doi.org/10.1006/jmaa.1993.1044>
124. H. M. Srivastava, G. Murugusundaramoorthy and S. Sivasubramanian, Hypergeometric functions in the parabolic starlike and uniformly convex domains, *Integral Transforms Spec. Funct.* 18 (2007), 511 – 520. <https://doi.org/10.1080/10652460701391324>
125. L. Spaček, Contribution à la théorie des fonctions univalentes, *Časopis Pro Pěstování Matematiky* 62(1932), 12 – 19.
126. Salah, Jamal Y. "A new subclass of univalent functions defined by the means of Jamal operator." *Far East Journal of Mathematical Sciences (FJMS)* Vol 108 (2018): 389-399.
127. Jamal Y. Salah On Riemann Hypothesis and Robin's Inequality. *International Journal of Scientific and Innovative Mathematical Research (IJSIMR)*. Volume (3) 4 (2015) 9-14.

128. Salah, Jamal. "Neighborhood of a certain family of multivalent functions with negative coefficients." *Int. J. Pure Appl. Math* 92.4 (2014): 591-597.
129. Salah, Jamal, and Maslina Darus. "A note on Starlike functions of order α associated with a fractional calculus operator involving Caputo's fractional." *J. Appl. comp Sc. Math* 5.1 (2011): 97-101.
130. J. Salah, Certain subclass of analytic functions associated with fractional calculus operator, *Trans. J. Math. Mech.*, 3 (2011), 35–42. Available from: <http://tjmm.edyopress.ro/journal/11030106.pdf>.
131. Salah, Jamal. "Some Remarks and Propositions on Riemann Hypothesis." *Mathematics and Statistics* 9.2 (2021): 159-165.
132. Salah, Jamal Y. Mohammad. "The consequence of the analytic continuity of Zeta function subject to an additional term and a justification of the location of the non-trivial zeros." *International Journal of Science and Research (IJSR)* 9.3 (2020): 1566-1569.
- 133.
134. A. Swaminathan, Certain sufficient conditions on Gaussian hypergeometric functions, *Journal of Inequalities in Pure and Applied Mathematics* 5(4) (2004), Article ID 83, 10 pages.
135. A. Wiman, Über die nullstellen der funktionen $E_\alpha(x)$, *Acta Math.* 29 (1905), 217-134.
136. Salah, Jamal Y. Mohammad. "An Alternative perspective to Riemann Hypothesis." *PSYCHOLOGY AND EDUCATION* 57.9 (2020): 1278-1281.
137. Salah, Jamal Y. "A note on the Hurwitz zeta function." *Far East Journal of Mathematical Sciences (FJMS)* 101.12 (2017): 2677-2683.
138. Jamal Y. Salah. Closed-to-Convex Criterion Associated to the Modified Caputo's fractional Calculus Derivative Operator. *Far East Journal of Mathematical Sciences (FJMS)*. Vol. 101, No. 1, pp. 55-59, 2017, DOI: 10.17654/MS101010055.
139. Jamal Y. Salah, A Note on Riemann Zeta Function, *IOSR Journal of Engineering (IOSRJEN)*, vol. 06, no. 02, pp. 07-16, February 2016, URL: [http://iosrjen.org/Papers/vol6_issue2%20\(part-3\)/B06230716.pdf](http://iosrjen.org/Papers/vol6_issue2%20(part-3)/B06230716.pdf)
140. Jamal Y. Salah, A Note on Gamma Function, *International Journal of Modern Sciences and Engineering Technology (IJMSET)*, vol. 2, no. 8, pp. 58-64, 2015.
141. Salah, Jamal, and S. Venkatesh. "Inequalities on the Theory of Univalent Functions." *Journal of Mathematics and System Science* 4.7 (2014).
142. T. R. Prabhakar, A single integral equation with a generalized Mittag – Leffler function in the kernel, *Yokohama Math. J.* 19 (1971), 7 – 15.
143. [Omar Jawabreh](#), [Ahmad Abdel Qader](#), [Jamal Salah](#), [Khaled Al Mashrafi](#), [Emad Al Dein AL Fahmawee](#), [Basel J. A. Ali](#), Fractional Calculus Analysis of Tourism Mathematical Model, *Progr. Fract. Differ. Appl.* Vol. 9, No. S1, pp. 1 -11 202) DOI: 10.18576/pfda/09S101 URL: <https://www.ajouronline.com/index.php/AJFAM/article/view/724>
144. Jamal Salah. Subordination and superordination involving certain fractional operator. *Asian Journal of Fuzzy and Applied Mathematics*, vol. 1, pp. 98-107, 2013. URL: <https://www.ajouronline.com/index.php/AJFAM/article/view/724>

145. Salah, Jamal Y. Mohammad. "Two Conditional proofs of Riemann Hypothesis." *International Journal of Sciences: Basic and Applied Research (IJSBAR)* 49.1 (2020): 74-83.
146. Salah, Jamal. "Fekete-Szego problems involving certain integral operator." *International Journal of Mathematics Trends and Technology-IJMTT* 7 (2014).